

AD _____

Award Number: DAMD17-02-1-0187

TITLE: Ethanol and Mesolimbic Serotonin/Dopamine Interactions
via 5HT-1B Receptors

PRINCIPAL INVESTIGATOR: Qingshan Yan, Ph.D.

CONTRACTING ORGANIZATION: University of Illinois at Chicago
Chicago, Illinois 60612-7227

REPORT DATE: March 2003

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20030930 038

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 074-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE March 2003	3. REPORT TYPE AND DATES COVERED Annual (15 Feb 02 - 14 Feb 03)		
4. TITLE AND SUBTITLE Ethanol and Mesolimbic Serotonin/Dopamine Interactions via 5HT-1B Receptors		5. FUNDING NUMBERS DAMD17-02-1-0187		
6. AUTHOR(S) : Qingshan Yan, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Illinois at Chicago Chicago, Illinois 60612-7227 E-Mail: QSY@uic.edu		8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012		10. SPONSORING / MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited			12b. DISTRIBUTION CODE	
13. Abstract (Maximum 200 Words) (abstract should contain no proprietary or confidential information) According to Statement of Work, Hypothesis 1 under Specific Aim 1 was tested during Year 1. To this end, dual-probe microdialysis was employed and the 5-HT1B receptor agonist CP 93129 used. The experiments were performed in freely-moving rats. One probe was in the ventral tegmental area (VTA) and the other in the ipsilateral nucleus accumbens (NACC). Artificial cerebrospinal fluid (ACSF) alone and ACSF with three concentrations of CP 93129 (20, 40, and 80 μ M) were infused respectively into the VTA of separate groups of rats for 60 min. Dialysates from the VTA and NACC were collected at 20-min intervals for determination of NACC DA and VTA GABA via HPLC systems. The results showed that administration of CP 93129 caused significant increases of extracellular DA concentrations in the NACC in a concentration-related manner. The maximum increases of NACC DA produced by 20, 40, and 80 μ M of CP 93129 were 163%, 181%, and 214% of baseline, respectively. The determination of GABA levels in the VTA is presently under way and has not been completed yet. The results suggest that activation of VTA 5-HT1B receptors by focally applied CP 93129 increases DA transmission in the ipsilateral NACC.				
14. SUBJECT TERMS: ethanol, dopamine, serotonin, 5HT-1B receptors, ventral tegmental area, nucleus accumbens			15. NUMBER OF PAGES 11	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

Table of Contents

Cover.....	1
SF 298.....	2
Introduction.....	4
Body.....	5-6
Key Research Accomplishments.....	7
Reportable Outcomes.....	8
Conclusions.....	9
References.....	10
Appendices.....	11

INTRODUCTION

The purpose of this project entitled "Ethanol and mesolimbic serotonin (5-HT) /dopamine (DA) interactions via 5-HT_{1B} receptors" is to investigate whether activation of 5-HT_{1B} receptors in the ventral tegmental area (VTA) facilitates DA transmission in the ipsilateral nucleus accumbens (NACC) and potentiates ethanol-induced increases in NACC DA by 5-HT_{1B} receptor-mediated GABA mechanisms. The scope of this project covers the following specific aims: (1) to determine the involvement of 5-HT_{1B} heteroreceptors on GABA terminals in the VTA in the modulation of GABA release in the VTA and DA release in the ipsilateral NACC, and its involvement in the neurochemical effect of acute ethanol in freely moving animals; (2) to compare the impact of 5-HT_{1B} receptor activation on DA transmission in the NACC and on ethanol's neurochemical effects between 5-HT_{1B} receptor knock-out mice and their counterparts wild-type mice; and (3) to determine the involvement of 5-HT_{1B} heteroreceptors on GABA terminals in the VTA in the modulation of DA and GABA releases in the VTA, and its involvement in the effect of ethanol in superfused VTA slices.

BODY

There are two hypotheses Under Specific aim 1: one is that activation of 5-HT_{1B} receptors in the VTA decreases GABA release in this area and increases DA transmission in the ipsilateral NACC and the other is that activation and blockade of VTA 5-HT_{1B} receptors potentiates and attenuates ethanol's effects on DA transmission in the ipsilateral NACC, respectively. According to Statement of Work, Hypothesis 1 was supposed to be completed at the end of Year 1. However, due to unexpected circumstances, the progress of the project has considerably lagged behind what had been originally proposed. This delay is caused mainly by search for a post-doctoral research associate. According to our university's policies, the search for that person could not be initiated until the funds were awarded. In addition, the search was a time-consuming process. As a result, the post-doctoral research associate did not participate in this project until August of 2002, a half year later than the proposed date. Therefore, a delay in hiring the post-doctoral research associate has caused a considerably delay in the progress of the project.

Hypothesis 1 consists of two parts of experiments. One is associated with the 5-HT_{1B} receptor agonist CP 93129. This part of experiments have almost been accomplished during Year 1. Due to the delay in the progress mentioned above, the second part of experiments will be postponed to Year 2. In the first part of experiments, the dual-probe microdialysis, a technically very difficult procedure, was used in awake and freely-moving adult Sprague-Dawley rats. One probe was inserted into the VTA and the other in the ipsilateral NACC. Both probed were perfused with artificial cerebrospinal fluid (ACSF). After basal DA release in the NACC was stable, ACSF alone and ACSF with three different concentrations of CP 93129 (20, 40, and 80 μ M), a selective 5-HT_{1B} receptor agonist^{1,2}, were infused respectively into the VTA of separate groups of rats for 60 min. The dialysates from both VTA and the NACC were collected at 20 min of intervals for determination of DA in the NACC and gamma-aminobutyric acid (GABA) in the VTA via HPLC systems. The basal DA levels in the accumbal extracellular fluids were (fmol/sample, mean \pm SEM): 33.15 \pm 4.75 (the control group, n = 7), 34.94 \pm 3.75 (the 20 μ M group, n = 6), 33.62 \pm 3.57 (the 40 μ M group n = 7), and 32.99 \pm 3.48 (the 80 μ M group, n = 7). As shown in the following figure, switching between syringes containing ACSF has no significant effects on the dialysate DA levels in the ipsilateral NACC. However, administration of CP 93129 through a dialysis probe into the VTA caused significant increases of extracellular DA concentrations in the NACC in a concentration-related manner. The maximum increases of NACC DA produced by 20, 40, and 80 μ M of CP 93129 were 163%, 181%, and 214% of baseline, respectively. The determination of GABA levels in the VTA is presently under way and has not been completed yet. Increased NACC DA after intra-tegmental CP 93129 is consistent with studies reported in the literature^{3,4}.




Fig. Effects of infusion of CP 93129 into the ventral tegmental area (VTA) on dopamine concentrations in the ipsilateral nucleus accumbens (NACC). CP 93129 (20, 40, and 80 μ M) was administered through a probe into the VTA indicated by the bar. Extracellular dopamine in the ipsilateral NACC was monitored by a second probe in this region. Results are mean \pm S.E.M. * $P < 0.05$, ** $P < 0.05$ as compared with the control (ACSF alone) group; + $P < 0.05$ as compared with the 40 μ M group (two-way ANOVA followed by Tukey test).

KEY RESEARCH ACCOMPLISHMENTS

1. We have developed a dual-probe microdialysis method that is technically very difficult.
2. We have established a dose-response of the 5-HT_{1B} receptor agonist CP 93129 on DA. We found that administration of CP 93129 into the VTA via a dialysis probe increased DA concentrations in the ipsilateral nucleus accumbens.

REPORTABLE OUTCOMES

None

CONCLUSIONS

The results obtained during Year 1 suggest that activation of VTA 5-HT1B receptors by focally applied CP 93129 increases DA transmission in the ipsilateral NACC. However, a firm conclusion could be drawn only after experiments with 5-HT1B receptor antagonists (i.e., the second part of experiments under Hypothesis 1) are completed.

REFERENCES

1. Chopin, P., Moret, C., Briley, M.: Neuropharmacology of 5-hydroxytryptamine_{1B/D} receptor ligands. *Pharmacol. Ther.* 62, 385-405, 1994.
2. Macor, J.E., Burkhart, C.A., Heym, J.H., Ives, J.L., Lebel, L.A., Newman, M.E., Nielsen, J.A., Ryan, K., Schulz, D.W., Torgersen, L.K., Koe, B.K.: 3-(1,2,5,6-Tetrahydropyrid-4-yl)pyrrolo[3,2-b]pyrid-5-one: A potent and selective serotonin (5-HT_{1B}) agonist and rotationally restricted phenolic analogue of 5-methoxy-3-(1,2,5,6-tetrahydropyrid-4-yl)indole. *J. Med. Chem.* 33, 2087-2093, 1990.
3. Parsons, L.H. O'Dell, L.E., Manzardo, A., Polis, I., Stouffer, D.: 5-HT_{1B} receptor modulation of mesolimbic dopamine neurotransmission: Effects of cocaine exposure. Program No. 9.2, Abstract, Society for Neuroscience, 2002.
4. Yan, Q.S., Yan, S.E.: Activation of 5-HT_{1B/1D} receptors in the mesolimbic dopamine system increases dopamine release from the nucleus accumbens. *Eur. J. Pharmacol.*, 418: 55-64, 2001.

APPENDICES

None